

NMR SPECTROSCOPY

*Prepared by D . Veera Sudarshan,
lecturer in chemistry,GDC,Porumamilla*

Nuclear Magnetic Resonance Spectroscopy

Introduction to NMR Spectroscopy

- Nuclear magnetic resonance spectroscopy is a powerful analytical technique used to characterize organic molecules by identifying carbon-hydrogen frameworks within molecules.
- Two common types of NMR spectroscopy are used to characterize organic structure: ^1H NMR is used to determine the type and number of H atoms in a molecule; ^{13}C NMR is used to determine the type of carbon atoms in the molecule.
- The source of energy in NMR is **radio waves** which have long wavelengths, and thus low energy and frequency.
- When low-energy radio waves interact with a molecule, they can change the **nuclear spins** of some elements, including ^1H and ^{13}C .

Nuclear Magnetic Resonance Spectroscopy


Introduction to NMR Spectroscopy

- When a charged particle such as a proton spins on its axis, it creates a **magnetic field**. Thus, the nucleus can be considered to be a tiny bar magnet.
- Normally, these tiny bar magnets are randomly oriented in space. However, in the presence of a magnetic field B_0 , they are oriented with or against this applied field. More nuclei are oriented with the applied field because this arrangement is lower in energy.
- The energy difference between these two states is very small (<0.1 cal).

A spinning proton
creates a magnetic field.




With no external magnetic field...



The nuclear magnets are
randomly oriented.

In a magnetic field...



The nuclear magnets are
oriented **with or against** B_0 .

Nuclear Magnetic Resonance Spectroscopy


Introduction to NMR Spectroscopy

- In a magnetic field, there are now two energy states for a proton: a lower energy state with the nucleus aligned in the same direction as B_0 , and a higher energy state in which the nucleus aligned against B_0 .
- When an external energy source ($h\nu$) that matches the energy difference (ΔE) between these two states is applied, energy is absorbed, causing the nucleus to “spin flip” from one orientation to another.
- The energy difference between these two nuclear spin states corresponds to the low frequency RF region of the electromagnetic spectrum.

Nuclear Magnetic Resonance Spectroscopy

Introduction to NMR Spectroscopy

- Thus, two variables characterize NMR: an applied magnetic field B_0 , the strength of which is measured in tesla (T), and the frequency ν of radiation used for resonance, measured in hertz (Hz), or megahertz (MHz)—(1 MHz = 10^6 Hz).



- A nucleus is in *resonance* when it absorbs RF radiation and “spin flips” to a higher energy state.

Nuclear Magnetic Resonance Spectroscopy

Introduction to NMR Spectroscopy

- The frequency needed for resonance and the applied magnetic field strength are proportionally related:

$$\nu \propto B_0$$


- The stronger the magnetic field, the larger the energy difference between the two nuclear spin states, and the higher the ν needed for resonance.
- NMR spectrometers are referred to as **300 MHz instruments**, **500 MHz instruments**, and so forth, depending on the frequency of the RF radiation used for resonance.
- These spectrometers use very powerful magnets to create a small but measurable energy difference between two possible spin states.

Nuclear Magnetic Resonance Spectroscopy

Introduction to NMR Spectroscopy

Figure 14.1

Schematic of an NMR spectrometer



An NMR spectrometer. The sample is dissolved in a solvent, usually CDCl_3 (deuteriochloroform), and placed in a magnetic field. A radiofrequency generator then irradiates the sample with a short pulse of radiation, causing resonance. When the nuclei fall back to their lower energy state, the detector measures the energy released, and a spectrum is recorded. The superconducting magnets in modern NMR spectrometers have coils that are cooled in liquid helium and conduct electricity with essentially no resistance.

Nuclear Magnetic Resonance Spectroscopy


Introduction to NMR Spectroscopy

- Protons in different environments absorb at slightly different frequencies, so they are distinguishable by NMR.
- The frequency at which a particular proton absorbs is determined by its **electronic environment**.
- The size of the magnetic field generated by the electrons around a proton determines where it absorbs.
- Modern NMR spectrometers use a constant magnetic field strength B_0 , and then a narrow range of frequencies is applied to achieve the resonance of all protons.
- **Only nuclei that contain odd mass numbers (such as ^1H , ^{13}C , ^{19}F and ^{31}P) or odd atomic numbers (such as ^2H and ^{14}N) give rise to NMR signals.**

Nuclear Magnetic Resonance Spectroscopy

^1H NMR—The Spectrum

- An NMR spectrum is a plot of the intensity of a peak against its chemical shift, measured in **parts per million (ppm)**.



Nuclear Magnetic Resonance Spectroscopy

^1H NMR—The Spectrum

- NMR absorptions generally appear as sharp peaks.
- Increasing chemical shift is plotted from left to right.
- Most protons absorb between 0-10 ppm.
- The terms “upfield” and “downfield” describe the relative location of peaks. Upfield means to the right. Downfield means to the left.
- NMR absorptions are measured relative to the position of a reference peak at 0 ppm on the δ scale due to tetramethylsilane (TMS). TMS is a volatile inert compound that gives a single peak upfield from typical NMR absorptions.

Nuclear Magnetic Resonance Spectroscopy

^1H NMR—The Spectrum

- The chemical shift of the x axis gives the position of an NMR signal, measured in ppm, according to the following equation:


$$\text{chemical shift (in ppm on the } \delta \text{ scale)} = \frac{\text{observed chemical shift (in Hz) downfield from TMS}}{\nu \text{ of the NMR spectrometer (in MHz)}}$$

- By reporting the NMR absorption as a fraction of the NMR operating frequency, we get units, ppm, that are independent of the spectrometer.
- Four different features of a ^1H NMR spectrum provide information about a compound's structure:
 - a. Number of signals
 - b. Position of signals
 - c. Intensity of signals.
 - d. Spin-spin splitting of signals.


Nuclear Magnetic Resonance Spectroscopy

^1H NMR—Number of Signals


- The number of NMR signals equals the number of different types of protons in a compound.
- Protons in different environments give different NMR signals.
- Equivalent protons give the same NMR signal.



All equivalent H's
1 NMR signal




2 types of H's
2 NMR signals




3 types of H's
3 NMR signals


- To determine equivalent protons in cycloalkanes and alkenes, always draw all bonds to hydrogen.




NOT



Draw



NOT




Nuclear Magnetic Resonance Spectroscopy


^1H NMR—Number of Signals

Figure 14.2


The number of ^1H NMR signals of some representative organic compounds




1 type of H
1 NMR signal



3 types of H's
3 NMR signals



2 types of H's
2 NMR signals




3 types of H's
3 NMR signals

Nuclear Magnetic Resonance Spectroscopy


^1H NMR—Number of Signals

- In comparing two H atoms on a ring or double bond, two protons are equivalent only if they are cis (or trans) to the same groups.




1,1-dichloroethylene

1 type of H
1 NMR signal



1-bromo-1-chloroethylene

2 types of H's
2 NMR signals




chloroethylene

3 types of H's
3 NMR signals

Nuclear Magnetic Resonance Spectroscopy


^1H NMR—Number of Signals

- Proton equivalency in cycloalkanes can be determined similarly.



cyclopropane

All H's are equivalent.
1 NMR signal




chlorocyclopropane

3 types of H's
3 NMR signals

Nuclear Magnetic Resonance Spectroscopy


^1H NMR—Enantiotopic and Diastereotopic Protons.



- When substitution of two H atoms by Z forms enantiomers, the two H atoms are equivalent and give a single NMR signal. These two H atoms are called *enantiotopic* protons.


Nuclear Magnetic Resonance Spectroscopy

^1H NMR—Enantiotopic and Diastereotopic Protons.




(*R*)-2-chlorobutane


H_a and H_b are **diastereotopic**.



substitution of H_a



substitution of H_b




diastereomers

- When substitution of two H atoms by Z forms diastereomers, the two H atoms are not equivalent, and give two NMR signals. These two H atoms are called *diastereotopic* protons.

Nuclear Magnetic Resonance Spectroscopy

^1H NMR—Position of Signals

- In the vicinity of the nucleus, the magnetic field generated by the circulating electron decreases the external magnetic field that the proton “feels”.
- Since the electron experiences a lower magnetic field strength, it needs a lower frequency to achieve resonance. Lower frequency is to the right in an NMR spectrum, toward a lower chemical shift, so **shielding** shifts the absorption upfield.



Nuclear Magnetic Resonance Spectroscopy

^1H NMR—Position of Signals

- The less shielded the nucleus becomes, the more of the applied magnetic field (B_0) it feels.
- This **deshielded** nucleus experiences a higher magnetic field strength, so it needs a higher frequency to achieve resonance.
- Higher frequency is to the left in an NMR spectrum, toward higher chemical shift—so deshielding shifts an absorption downfield.
- Protons near electronegative atoms are deshielded, so they absorb downfield.

Nuclear Magnetic Resonance Spectroscopy


^1H NMR—Position of Signals

Figure 14.3

How chemical shift is affected by electron density around a nucleus

a. Shielding effects


- An electron shields the nucleus.
- The absorption shifts *upfield*.



← Increasing chemical shift
Increasing ν

b. Deshielding effects

- Decreased electron density deshields a nucleus.
- The absorption shifts *downfield*.




← Increasing chemical shift
Increasing ν

Nuclear Magnetic Resonance Spectroscopy

^1H NMR—Position of Signals


Figure 14.4

Shielding and deshielding effects




Nuclear Magnetic Resonance Spectroscopy


^1H NMR—Position of Signals



- The H_b protons are **deshielded** because they are closer to the electronegative Cl atom, so they absorb **downfield** from H_a .



- Because F is more electronegative than Br, the H_b protons are more **deshielded** than the H_a protons and absorb farther **downfield**.



- The larger number of electronegative Cl atoms (two versus one) **deshields** H_b more than H_a , so it absorbs **downfield** from H_a .

Nuclear Magnetic Resonance Spectroscopy








^1H NMR—Chemical Shift Values

- Protons in a given environment absorb in a predictable region in an NMR spectrum.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

Table 14.1


Characteristic Chemical Shifts of Common Types of Protons

Type of proton	Chemical shift (ppm)	Type of proton	Chemical shift (ppm)
 sp^3	0.9–2	 sp^2	4.5–6
<ul style="list-style-type: none"> RCH_3 R_2CH_2 R_3CH 	~0.9 ~1.3 ~1.7		6.5–8
 $\text{Z} = \text{C}, \text{O}, \text{N}$	1.5–2.5		9–10
$-\text{C}\equiv\text{C}-\text{H}$	~2.5		10–12
 sp^3 $\text{Z} = \text{N}, \text{O}, \text{X}$	2.5–4	$\text{RO}-\text{H}$ or $\text{R}-\text{N}-\text{H}$	1–5

Nuclear Magnetic Resonance Spectroscopy

^1H NMR—Chemical Shift Values


- The chemical shift of a C—H bond increases with increasing alkyl substitution.



Nuclear Magnetic Resonance Spectroscopy

^1H NMR—Chemical Shift Values


- In a magnetic field, the six π electrons in benzene circulate around the ring creating a **ring current**.
- The magnetic field induced by these moving electrons reinforces the applied magnetic field in the vicinity of the protons.
- The protons thus feel a stronger magnetic field and a higher frequency is needed for resonance. Thus they are deshielded and absorb downfield.



Nuclear Magnetic Resonance Spectroscopy

^1H NMR—Chemical Shift Values


- In a magnetic field, the loosely held π electrons of the double bond create a magnetic field that reinforces the applied field in the vicinity of the protons.
- The protons now feel a stronger magnetic field, and require a higher frequency for resonance. Thus the protons are deshielded and the absorption is downfield.



Nuclear Magnetic Resonance Spectroscopy

^1H NMR—Chemical Shift Values

- In a magnetic field, the π electrons of a carbon-carbon triple bond are induced to circulate, but in this case the induced magnetic field opposes the applied magnetic field (B_0).
- Thus, the proton feels a weaker magnetic field, so a lower frequency is needed for resonance. The nucleus is shielded and the absorption is upfield.





Nuclear Magnetic Resonance Spectroscopy

^1H NMR—Chemical Shift Values

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

Table 14.2

Effect of π Electrons on Chemical Shift Values


Proton type	Effect	Chemical shift (ppm)
	highly deshielded	6.5–8
	deshielded	4.5–6
$\text{—C}\equiv\text{C—H}$	shielded	~2.5

Nuclear Magnetic Resonance Spectroscopy

^1H NMR—Chemical Shift Values)

Figure 14.5

Regions in the ^1H NMR spectrum



- Shielded protons absorb at lower chemical shift (to the right).
- Deshielded protons absorb at higher chemical shift (to the left).


Nuclear Magnetic Resonance Spectroscopy

^1H NMR—Intensity of Signals

- The area under an NMR signal is proportional to the number of absorbing protons.
- An NMR spectrometer automatically **integrates** the area under the peaks, and prints out a stepped curve (**integral**) on the spectrum.
- The height of each step is proportional to the area under the peak, which in turn is proportional to the number of absorbing protons.
- Modern NMR spectrometers automatically calculate and plot the value of each integral in arbitrary units.
- The ratio of integrals to one another gives the ratio of absorbing protons in a spectrum. Note that this gives a ratio, and not the absolute number, of absorbing protons.

Nuclear Magnetic Resonance Spectroscopy


^1H NMR—Intensity of Signals



How To

Determine the Number of Protons Giving Rise to an NMR Signal

Example A compound of molecular formula $C_9H_{10}O_2$ gives the following integrated 1H NMR spectrum. How many protons give rise to each signal?



Step [1] Determine the number of integration units per proton by dividing the total number of integration units by the total number of protons.

- Total number of integration units: $54 + 23 + 33 = 110$ units
- Total number of protons = 10
- Divide: $110 \text{ units}/10 \text{ protons} = \mathbf{11 \text{ units per proton}}$

Step [2] Determine the number of protons giving rise to each signal.


- To determine the number of H atoms giving rise to each signal, divide each integration value by the answer of Step [1] and round to the nearest whole number.

$$\text{Answer: } \begin{array}{l} \text{Signal [A]:} \\ \frac{54}{11} = 4.9 \approx \mathbf{5 H} \end{array} \quad \begin{array}{l} \text{Signal [B]:} \\ \frac{23}{11} = 2.1 \approx \mathbf{2 H} \end{array} \quad \begin{array}{l} \text{Signal [C]:} \\ \frac{33}{11} = \mathbf{3 H} \end{array}$$

Nuclear Magnetic Resonance Spectroscopy

^1H NMR—Spin-Spin Splitting

- Consider the spectrum below:



Nuclear Magnetic Resonance Spectroscopy

^1H NMR—Spin-Spin Splitting

- **Spin-spin splitting** occurs only between **nonequivalent protons** on the same carbon or adjacent carbons.

Let us consider how the **doublet** due to the CH_2 group on $\text{BrCH}_2\text{CHBr}_2$ occurs:

- When placed in an applied magnetic field, (B_0), the adjacent proton (CHBr_2) can be aligned with (\uparrow) or against (\downarrow) B_0 .
- Thus, the absorbing CH_2 protons feel two slightly different magnetic fields—one slightly larger than B_0 , and one slightly smaller than B_0 .
- Since the absorbing protons feel two different magnetic fields, they absorb at two different frequencies in the NMR spectrum, thus splitting a single absorption into a doublet.


Nuclear Magnetic Resonance Spectroscopy

^1H NMR—Spin-Spin Splitting


The frequency difference, measured in Hz between two peaks of the doublet is called the **coupling constant, J** .

How a doublet arises


With no adjacent H's:
The absorbing H's feel only one magnetic field.




The NMR signal is a **single peak**.



With one adjacent H:
The absorbing H's feel two different fields, so they absorb at two different frequencies.



two different magnetic fields



1:1

The NMR signal is split into a **doublet**.

- One adjacent proton splits an NMR signal into a doublet.

Nuclear Magnetic Resonance Spectroscopy

^1H NMR—Spin-Spin Splitting

Let us now consider how a triplet arises:



- When placed in an applied magnetic field (B_0), the adjacent protons H_a and H_b can each be aligned with (\uparrow) or against (\downarrow) B_0 .
- Thus, the absorbing proton feels three slightly different magnetic fields—one slightly larger than B_0 , one slightly smaller than B_0 , and one the same strength as B_0 .


Nuclear Magnetic Resonance Spectroscopy

^1H NMR—Spin-Spin Splitting

- Because the absorbing proton feels three different magnetic fields, it absorbs at three different frequencies in the NMR spectrum, thus splitting a single absorption into a triplet.
- Because there are two different ways to align one proton with B_0 , and one proton against B_0 —that is, $\uparrow_a\downarrow_b$ and $\downarrow_a\uparrow_b$ —the middle peak of the triplet is twice as intense as the two outer peaks, making the ratio of the areas under the three peaks 1:2:1.
- Two adjacent protons split an NMR signal into a triplet.
- When two protons split each other, they are said to be coupled.
- The spacing between peaks in a split NMR signal, measured by the J value, is equal for coupled protons.

Nuclear Magnetic Resonance Spectroscopy

^1H NMR—Spin-Spin Splitting




Nuclear Magnetic Resonance Spectroscopy

^1H NMR—Spin-Spin Splitting



Three general rules describe the splitting patterns commonly seen in the ^1H NMR spectra of organic compounds.

1. Equivalent protons do not split each other's signals.
2. A set of n nonequivalent protons splits the signal of a nearby proton into $n + 1$ peaks.
3. Splitting is observed for nonequivalent protons on the same carbon or adjacent carbons.

If H_a and H_b are not equivalent, splitting is observed when:



H_a and H_b are on the **same** carbon.




H_a and H_b are on **adjacent** carbons.

Nuclear Magnetic Resonance Spectroscopy

^1H NMR—Spin-Spin Splitting


Splitting is not generally observed between protons separated by more than three σ bonds.



2-butanone

H_a and H_b are separated by four σ bonds.

no splitting between H_a and H_b



ethyl methyl ether

H_a and H_b are separated by four σ bonds.

no splitting between H_a and H_b











Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

Table 14.3

Names for a Given Number of Peaks in an NMR Signal

Number of peaks	Name	Number of peaks	Name
1	singlet	5	quintet
2	doublet	6	sextet
3	triplet	7	septet
4	quartet	> 7	multiplet

Table 14.4**Common Splitting Patterns Observed in ^1H NMR**

Example	Pattern	Analysis (H_a and H_b are not equivalent.)
[1] 		<ul style="list-style-type: none"> • H_a: one adjacent H_b proton \dashrightarrow two peaks \dashrightarrow a doublet • H_b: one adjacent H_a proton \dashrightarrow two peaks \dashrightarrow a doublet
[2] 		<ul style="list-style-type: none"> • H_a: two adjacent H_b protons \dashrightarrow three peaks \dashrightarrow a triplet • H_b: one adjacent H_a proton \dashrightarrow two peaks \dashrightarrow a doublet
[3] 		<ul style="list-style-type: none"> • H_a: two adjacent H_b protons \dashrightarrow three peaks \dashrightarrow a triplet • H_b: two adjacent H_a protons \dashrightarrow three peaks \dashrightarrow a triplet
[4] 		<ul style="list-style-type: none"> • H_a: three adjacent H_b protons \dashrightarrow four peaks \dashrightarrow a quartet* • H_b: two adjacent H_a protons \dashrightarrow three peaks \dashrightarrow a triplet
[5] 		<ul style="list-style-type: none"> • H_a: three adjacent H_b protons \dashrightarrow four peaks \dashrightarrow a quartet* • H_b: one adjacent H_a proton \dashrightarrow two peaks \dashrightarrow a doublet

*The relative area under the peaks of a quartet is 1:3:3:1.


Nuclear Magnetic Resonance Spectroscopy

^1H NMR—Spin-Spin Splitting

Whenever two (or three) different sets of adjacent protons are equivalent to each other, use the $n + 1$ rule to determine the splitting pattern.

Figure 14.6

The ^1H NMR spectrum of 2-bromopropane, $[(\text{CH}_3)_2\text{CHBr}]$



Nuclear Magnetic Resonance Spectroscopy


^1H NMR—Spin-Spin Splitting

Now consider the spectrum of 1-bromopropane. Since H_a and H_c are not equivalent to each other, we cannot merely add them together and use the $n + 1$ rule.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

Figure 14.7

The ^1H NMR spectrum of 1-bromopropane, $\text{CH}_3\text{CH}_2\text{CH}_2\text{Br}$




- H_a and H_c are both triplets.
- H_b is split into 12 peaks, labeled as a multiplet. Fewer peaks are seen because some peaks overlap.

Nuclear Magnetic Resonance Spectroscopy

^1H NMR—Spin-Spin Splitting

When two sets of adjacent protons are different from each other (n protons on one adjacent carbon and m protons on the other), the number of peaks in an NMR signal = $(n + 1)(m + 1)$.



Three H_a protons split the H_b signal into $3 + 1 = 4$ peaks.

Two H_c protons further split the H_b signal into $2 + 1 = 3$ peaks.




Figure 14.8


A splitting diagram for the H_b protons in 1-bromopropane

- The H_b signal is split into 12 peaks, a quartet of triplets. The number of peaks actually seen for the signal depends on the relative size of the coupling constants, J_{ab} and J_{bc} . When $J_{ab} \gg J_{bc}$, as drawn in this diagram, all 12 lines of the pattern are visible. When J_{ab} and J_{bc} are similar in magnitude, peaks overlap and fewer lines are observed.

Nuclear Magnetic Resonance Spectroscopy

^1H NMR—Spin-Spin Splitting

- Protons on carbon-carbon double bonds often give characteristic splitting patterns.
- A disubstituted double bond can have two geminal protons, two cis protons, or two trans protons.
- When these protons are different, each proton splits the NMR signal of the other so that each proton appears as a doublet.
- The magnitude of the coupling constant J for these doublets depends on the arrangement of hydrogen atoms.




characteristic coupling constants for three types of disubstituted alkenes

Nuclear Magnetic Resonance Spectroscopy

^1H NMR—Spin-Spin Splitting

Figure 14.9

^1H NMR spectra for the alkenyl protons of (*E*)- and (*Z*)-3-chloropropenoic acid



- Although both (*E*)- and (*Z*)-3-chloropropenoic acid show two doublets in their ^1H NMR spectra for their alkenyl protons, $J_{\text{trans}} > J_{\text{cis}}$.


Nuclear Magnetic Resonance Spectroscopy

^1H NMR—Spin-Spin Splitting

Figure 14.10

The ^1H NMR spectrum of vinyl acetate ($\text{CH}_2=\text{CHOCOCH}_3$)

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.




Nuclear Magnetic Resonance Spectroscopy

^1H NMR—Spin-Spin Splitting

Splitting diagrams for the alkenyl protons in vinyl acetate are shown below. Note that each pattern is different in appearance because the magnitude of the coupling constants forming them is different.


Figure 14.11

Splitting diagram for the alkenyl protons in vinyl acetate ($\text{CH}_2=\text{CHOCH}_3$)



$J_{bc} = 1.2$ Hz (geminal)
 $J_{cd} = 6.5$ Hz (cis)
 $J_{bd} = 14$ Hz (trans)

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



One nearby H splits the signal into a doublet.

The second nearby proton splits the doublet into a doublet of doublets.


Nuclear Magnetic Resonance Spectroscopy

^1H NMR—OH Protons

- Under usual conditions, an OH proton does not split the NMR signal of adjacent protons.
- The signal due to an OH proton is not split by adjacent protons.

Figure 14.12

The ^1H spectrum of ethanol ($\text{CH}_3\text{CH}_2\text{OH}$)



Nuclear Magnetic Resonance Spectroscopy


^1H NMR—OH Protons

- Ethanol ($\text{CH}_3\text{CH}_2\text{OH}$) has three different types of protons, so there are three signals in its NMR spectrum.
- The H_a signal is split by the two H_b protons into three peaks (a triplet).
- The H_b signal is split only by the three H_a protons into four peaks, a quartet. The adjacent OH proton does not split the signal due to H_b .
- H_c is a singlet because OH protons are not split by adjacent protons.
- Protons on electronegative atoms rapidly exchange between molecules in the presence of trace amounts of acid or base. Thus, the CH_2 group of ethanol never “feels” the presence of the OH proton, because the OH proton is rapidly moving from one molecule to another.
- This phenomenon usually occurs with NH and OH protons.

Nuclear Magnetic Resonance Spectroscopy

^1H NMR—Cyclohexane Conformers

- Recall that cyclohexane conformers interconvert by ring flipping.
- Because the ring flipping is very rapid at room temperature, an NMR spectrum records an average of all conformers that interconvert.
- Thus, even though each cyclohexane carbon has two different types of hydrogens—one axial and one equatorial—the two chair forms of cyclohexane rapidly interconvert them, and an NMR spectrum shows a single signal for the average environment that it “sees”.




Nuclear Magnetic Resonance Spectroscopy

^1H NMR—Protons on Benzene Rings







- Benzene has six equivalent deshielded protons and exhibits a single peak in its ^1H NMR spectrum at 7.27 ppm.
- Monosubstituted benzenes contain five deshielded protons that are no longer equivalent, and the appearance of these signals is highly variable, depending on the identity of Z.

Figure 14.13 The 6.5–8 ppm region of the ^1H NMR spectrum of three benzene derivatives

A benzene ring with one substituent Z



A monosubstituted benzene ring has three different types of H atoms: H_a , H_b , and H_c .




- The appearance of the signals in the 6.5–8 ppm region of the ^1H NMR spectrum depends on the identity of Z in $\text{C}_6\text{H}_5\text{Z}$.

Nuclear Magnetic Resonance Spectroscopy

^1H NMR—Structure Determination

How To Use ^1H NMR Data to Determine a Structure

Example Using its ^1H NMR spectrum, determine the structure of an unknown compound X that has molecular formula $\text{C}_4\text{H}_8\text{O}_2$ and contains a $\text{C}=\text{O}$ absorption in its IR spectrum.



Step [1] Determine the number of different kinds of protons.

- The number of NMR signals equals the number of different types of protons.
- This molecule has three NMR signals ([A], [B], and [C]) and therefore **three** types of protons (H_a , H_b , and H_c).


Nuclear Magnetic Resonance Spectroscopy

^1H NMR—Structure Determination

How To, continued . . .

Step [2] Use the integration data to determine the number of H atoms giving rise to each signal (Section 14.5).

- Total number of integration units: $14 + 11 + 15 = 40$ units
- Total number of protons = 8
- Divide: $40 \text{ units} / 8 \text{ protons} = 5 \text{ units per proton}$
- Then, divide each integration value by this answer (5 units per proton) and round to the nearest whole number.




Nuclear Magnetic Resonance Spectroscopy

^1H NMR—Structure Determination


How To, continued . . .

Step [3] Use individual splitting patterns to determine what carbon atoms are bonded to each other.

- Start with the singlets. Signal [C] is due to a CH_3 group with no adjacent nonequivalent H atoms. Possible structures include:



- Because signal [A] is a **triplet**, there must be **2 H's** (CH_2 group) on the adjacent carbon.
- Because signal [B] is a **quartet**, there must be **3 H's** (CH_3 group) on the adjacent carbon.
- This information suggests that **X** has an **ethyl** group $\text{---} \rightarrow \text{CH}_3\text{CH}_2\text{---}$.



To summarize, **X** contains $\text{CH}_3\text{---}$, $\text{CH}_3\text{CH}_2\text{---}$, and $\text{C}=\text{O}$ (from the IR). Comparing these atoms with the molecular formula shows that one O atom is missing. Because O atoms do not absorb in a ^1H NMR spectrum, their presence can only be inferred by examining the chemical shift of protons near them. O atoms are more electronegative than C, thus deshielding nearby protons, and shifting their absorption downfield.


Nuclear Magnetic Resonance Spectroscopy

^1H NMR—Structure Determination

How To, continued . . .

Step [4] Use chemical shift data to complete the structure.

- Put the structure together in a manner that preserves the splitting data and is consistent with the reported chemical shifts.
- In this example, two isomeric structures (**A** and **B**) are possible for **X** considering the splitting data only:




- Chemical shift information distinguishes the two possibilities. The electronegative O atom deshields adjacent H's, shifting them downfield between 3 and 4 ppm. If **A** is the correct structure, the singlet due to the CH_3 group (H_c) should occur downfield, whereas if **B** is the correct structure, the quartet due to the CH_2 group (H_b) should occur downfield.
- Because the NMR of **X** has a singlet (not a quartet) at 3.7, **A is the correct structure**.

Nuclear Magnetic Resonance Spectroscopy

^{13}C NMR

^{13}C Spectra are easier to analyze than ^1H spectra because the signals are not split. Each type of carbon atom appears as a single peak.



Nuclear Magnetic Resonance Spectroscopy


^{13}C NMR

- The lack of splitting in a ^{13}C spectrum is a consequence of the low natural abundance of ^{13}C .
- Recall that splitting occurs when two NMR active nuclei—like two protons—are close to each other. Because of the low natural abundance of ^{13}C nuclei (1.1%), the chance of two ^{13}C nuclei being bonded to each other is very small (0.01%), and so no carbon-carbon splitting is observed.
- A ^{13}C NMR signal can also be split by nearby protons. This ^1H - ^{13}C splitting is usually eliminated from the spectrum by using an instrumental technique that decouples the proton-carbon interactions, so that every peak in a ^{13}C NMR spectrum appears as a singlet.
- The two features of a ^{13}C NMR spectrum that provide the most structural information are the number of signals observed and the chemical shifts of those signals.

Nuclear Magnetic Resonance Spectroscopy

^{13}C NMR—Number of Signals


- The number of signals in a ^{13}C spectrum gives the number of different types of carbon atoms in a molecule.
- Because ^{13}C NMR signals are not split, the number of signals equals the number of lines in the ^{13}C spectrum.
- In contrast to the ^1H NMR situation, peak intensity is not proportional to the number of absorbing carbons, so ^{13}C NMR signals are not integrated.



dimethyl ether


1 ^{13}C NMR signal

Both C's are equivalent.



chloroethane

2 ^{13}C NMR signals



methyl acetate

3 ^{13}C NMR signals

Nuclear Magnetic Resonance Spectroscopy







^{13}C NMR—Position of Signals

- In contrast to the small range of chemical shifts in ^1H NMR (1-10 ppm usually), ^{13}C NMR absorptions occur over a much broader range (0-220 ppm).
- The chemical shifts of carbon atoms in ^{13}C NMR depend on the same effects as the chemical shifts of protons in ^1H NMR.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

Table 14.5

Common ^{13}C Chemical Shift Values

Type of carbon	Chemical shift (ppm)	Type of carbon	Chemical shift (ppm)
	5–45		100–140
 Z = N, O, X	30–80		120–150
	65–100		160–210

Nuclear Magnetic Resonance Spectroscopy


^{13}C NMR—Number of Signals

Figure 14.14

Representative ^{13}C NMR spectra

- The three types of C's in 1-propanol—identified as C_a , C_b , and C_c —give rise to three ^{13}C NMR signals.
- Deshielding increases with increasing proximity to the electronegative O atom, and the absorption shifts downfield; thus, in order of increasing chemical shift: $\text{C}_a < \text{C}_b < \text{C}_c$.


a. 1-Propanol



Nuclear Magnetic Resonance Spectroscopy

^{13}C NMR—Number of Signals

b. Methyl acetate



- The three types of C's in methyl acetate—identified as C_a , C_b , and C_c —give rise to three ^{13}C NMR signals.
- The carbonyl carbon (C_b) is highly deshielded, so it absorbs farthest downfield.
- C_a , an sp^3 hybridized C that is not bonded to an O atom, is the most shielded, and so it absorbs farthest upfield.
- Thus, in order of increasing chemical shift: $\text{C}_a < \text{C}_c < \text{C}_b$.

Nuclear Magnetic Resonance Spectroscopy

Figure 14.15

The MRI image of the lower back

- A:** Spinal cord compression from a herniated disc
- B:** Spinal cord (would not be visualized with conventional X-rays)

An MRI instrument is especially useful for visualizing soft tissue. In 2002, 60 million MRI procedures were performed. The 2003 Nobel Prize in Physiology or Medicine was awarded to chemist Paul C. Lauterbur and physicist Sir Peter Mansfield for their contributions in developing magnetic resonance imaging.

